

Activation of viral defense signaling in cancer

Maria Gonzalez-Cao, Niki Karachaliou, Mariacarmela Santarpia, Santiago Viteri, Andreas Meyerhans and Rafael Rosell

Abstract: A coordinated action of innate and adaptive immune responses is required to efficiently combat a microbial infection. It has now become clear that cancer therapies also largely benefit when both arms of the immune response are engaged. In this review, we will briefly describe the current knowledge of innate immunity and how this can be utilized to prime tumors for a better response to immune checkpoint inhibitors. Comments on compounds in development and ongoing clinical trials will be provided.

Keywords: cancer immunotherapy, innate immunity, TLRs, viral defense signaling

Received: 1 November 2017; revised manuscript accepted: 17 July 2018.

Introduction

Recent advances in cancer immunotherapy have been achieved with antibodies that inhibit immune checkpoint receptors on immune cells and tumor cells, mainly programmed-cell death 1 (PD-1), its ligand PD-L1 and cytotoxic T lymphocyte-associated molecule-4 (CTLA-4).¹ However, most cancer patients do not respond to single checkpoint inhibition.²⁻⁴ Among the reasons is the lack of tumor infiltration by cytotoxic CD8⁺ T-cells.⁵ Non-T-cell-infiltrated tumors ('cold tumors') probably require the combination of checkpoint inhibitors with other therapies designed to attract these effector cells into the tumor microenvironment.

The immune system has two arms, (i) the innate immune arm that is rapidly activated after an appropriate stimulus but lacks antigen-specificity and memory, and (ii) the adaptive immune response that requires time to appear but is antigen-specific and long-lasting.⁶ Both arms of the immune system are intimately linked such that the innate arm provides the conditions for an efficient activation of the adaptive response (Figure 1).

Elements of the innate defense system are physical barriers, soluble factors such as complement proteins, interferons (IFNs) and IFN-stimulated proteins, and immune cells such as dendritic cells, macrophages, neutrophils and natural killer

(NK) cells. The adaptive arm of the immune system comprises T and B lymphocyte subpopulations that recognize pathogens in an antigen-specific way *via* divergent T-cell receptors and B-cell receptors, respectively. Among the latter are CD4⁺ T-helper and T-regulatory cells (Tregs), cytotoxic CD8⁺ T-cells, B-cells and antibody-producing plasma cells.⁶

Several virus infections including *Papilloma* viruses, Merkel cell polyomavirus, and hepatitis B and C viruses can induce tumors in humans. Interestingly, such tumors that are linked to virus infections seem to respond better to checkpoint inhibitors than tumors that are not virus-linked.^{7,8} The reason for this seems related to a more activated innate immune response.⁹⁻¹⁶ Thus, like in virus infections, the innate immune system probably provides a better microenvironment for the development of a potent specific antitumor response.^{17,18}

Type I IFNs are key regulatory elements in this aspect. They are produced for example when virus components or cell-derived damage-associated molecular patterns (DAMPs) bind and activate pattern recognition receptors (PRRs).¹⁹ The secreted IFNs can then activate dendritic cells (DCs) in tumor-draining lymph nodes and enhance the cross-presentation of tumor-associated antigens to CD8⁺ T-cells,²⁰ which

Ther Adv Med Oncol

2018, Vol. 10: 1-12

DOI: 10.1177/
1758835918793105

© The Author(s), 2018.
Article reuse guidelines:
sagepub.com/journals-
permissions

Correspondence to:

Gonzalez-Cao Maria
Rosell Oncology Institute
(IOR), Dexeus University
Hospital, Quironsalud
Group, C/ Sabino Arana,
5, Barcelona 08028, Spain
[mgonzalezcao@
oncorosell.com](mailto:mgonzalezcao@oncorosell.com)

Niki Karachaliou
Rosell Oncology
Institute (IOR), Sagrat
Cor University Hospital,
Quironsalud Group,
Barcelona, Spain

Mariacarmela Santarpia
Medical Oncology Unit,
Department of Human
Pathology 'G. Barresi',
University of Messina,
Messina, Italy

Santiago Viteri
Rosell Oncology Institute
(IOR), Dexeus University
Hospital, Quironsalud
Group, Barcelona, Spain
Rosell Oncology Institute
(IOR), Teknon Medical
Center, Quironsalud
Group, Barcelona, Spain

Andreas Meyerhans
Infection Biology
Laboratory, Department
of Experimental and
Health Sciences (DCEXS),
Universitat Pompeu Fabra,
Barcelona, Spain
Institució Catalana de
Recerca i Estudis Avançats
(ICREA), Barcelona, Spain

Rafael Rosell
Rosell Oncology Institute
(IOR), Dexeus University
Hospital, Quironsalud
Group, Barcelona, Spain
Rosell Oncology
Institute (IOR), Sagrat
Cor University Hospital,
Quironsalud Group,
Barcelona, Spain
Catalan Institute of
Oncology, Germans Trias i
Pujol University Hospital,
Badalona, Spain




Figure 1. From stimulating innate immune responses to an adaptive antitumor response. A schematic view is given. In the tumor microenvironment, activation of PRRs is achieved by the recognition of DAMPs released by dying tumor cells and by drugs such as TLR agonists, STING agonists, DNA demethylating agents and treatments based on the use of modified oncolytic viruses. The activation of the innate immune response leads to an efficient priming by dendritic cells of T-cells in lymph nodes and to the infiltration by tumor-specific T-cells into the tumor. Details are described in the text.

DAMP, danger-associated molecular pattern; ISG, interferon-stimulated gene; PRR, pattern recognition receptor; TLR, Toll-like receptor; STING, stimulator of interferon genes complex.

subsequently may lead to tumor-specific CD8⁺ T-cell expansion and tumor destruction.

A detailed understanding of the innate immune response against viruses may provide opportunities for developing more efficient treatments in the field of cancer immunotherapy. Indeed, the combination of checkpoint inhibitors with agents that trigger the innate immune response enhances their antitumor effect. Here, we will review the recently described means of activating innate immune responses to improve immunotherapy for cancer patients. Novel strategies that activate directly or indirectly PRRs will be commented on.

Pathogen recognition receptors (PRRs)

The innate immune response initiates with the recognition of foreign nucleic acids or other molecules in host cells by PRRs. PRRs recognize molecules derived from pathogen-associated molecular patterns, as well as DAMPs released from endogenous tissues that have suffered some damage.²¹ The understanding of the immunostimulatory as

well as pro- or antitumoral function of PRRs is necessary to exploit them for enhancing cancer immunotherapy.

Several different subtypes of PRRs are described today: Toll-like receptors (TLRs), NOD-like receptors (NLRs), c-type lectin receptors (CLRs), cytosol dsDNA sensors (CDSs) and retinol acid inducible gene 1 (RIG-1)-like receptors (RLRs).^{22,23} PRRs are classified according to their cellular location. They are located in cell membranes, such as TLRs, or in the cytoplasm like NLRs, CLRs, CDSs and RIG-1-like receptors (RLRs).²¹ PRR activation induces the production of type I IFNs (mainly IFN- α proteins and IFN- β).²² Subsequently, type I IFNs control the transcription of genes that are restricting viral infections (so-called ‘virus restriction factors’). In addition, type I IFNs activate NK cells, promote antigen presentation²⁴ and participate in the differentiation of specific CD8⁺ cytotoxic T lymphocytes (CTLs). Finally, type I IFNs have antiproliferative functions that are through TP53 gene induction.^{25,26}




Figure 2. Main subtypes of PRRs as targets for cancer treatment: Toll-like receptors (TLR3, TLR7/8, TLR9), cytosol dsDNA sensors (cGAS/IFI that activate STING) and the retinol acid inducible gene RIG-1 like receptors (RIG1 and MDA5). Activation of these receptors, following detection of nucleic acids from virus, induces production of type I IFNs (IFN- α and IFN- β). dsRNA (from HERVs re-expressed after treatment with azacytidine or from exogenous infection) activates TLR3, MDA5 and RIG1; ssRNA activates TLR7/8; DNA (from pathogens or from tumor cells) activates STING.

HERV, human endogenous retrovirus; IFN, interferon; IL, interleukin; IRF-3, IFN-regulatory factor-3; NF κ B, nuclear factor kappa B; PRR, pathogen recognition receptor; ssRNA, single stranded RNA; STING, stimulator of interferon genes complex; TBK1, TANK-binding kinase 1; TNF, tumor necrosis factor; TRAF6, TNF receptor-associated factor 6; TRIF, Toll/IL-1 receptor domain-containing adaptor inducing IFN- β .

TLRs constitute a receptor family that is mainly expressed on macrophages and DCs. In humans, the family has 11 members²⁷ located on the extracellular membrane (TLR 1, 2, 4, 5, 6 and 11) or in the intracellular counterpart of endosomes (TLR 2, 3, 7, 8, 9 and 10). Cell membrane-bound TLRs recognize glycoproteins, while endosome-placed TLRs respond to nucleic acid molecules, in particular viral RNA.^{27–29}

TLR4 was the first TLR identified. Activation of TLR4 signaling is preceded by binding of lipopolysaccharides produced by Gram-negative bacteria.^{30,31} In cancer, TLR4 activation has a dual role. Although its upregulation is associated with chemoresistance,³² metastasis and immunosuppression³³ in several tumor types, TLR4 activation has also an anticancer effect. While TLR4 antagonists could help reduce metastasis, TLR4 agonists have been shown to induce antitumor immunity in patients and models of cancer.

Several TLR4 agonists, such as OM-174,³⁴ or the *Streptococcus*-derived agent OK-432,^{35,36} Coley toxin (a mixture of killed *Streptococcus pyogenes* and *Serratiamarcescens* bacteria) and *Bacillus Calmette-Guerin*, have antitumoral effects.^{36–38}

Double-stranded RNA (dsRNA) is detected *via* TLR3 and RLRs (RIG1 and MDA5). Among immune cells, myeloid DCs and macrophages express TLR3. TLR3 is also expressed in fibroblasts and hepatocytes. When activated, TLR3, through TIR-domain-containing adapter-inducing interferon- β (TRIF), tumor necrosis factor (TNF) receptor-associated factor 6 (TRAF6) and tankyrase 1 (TANK1), activates the transcription factors interferon response factor 3 (IRF-3) and nuclear factor kappa B (NF- κ B). This then leads to the expression of type I IFNs, mainly IFN- β ³⁹ (Figure 2). RLRs use protein adaptor mitochondrial antiviral signaling to activate IRF-3, IRF-7 and NF- κ B⁴⁰ (Figure 2). RLRs

Table 1. Cancer clinical trials in progress with drugs targeting innate immune response.

Drug class	Drug	Target	In combination with	Tumor type	NCT number	
PRR	MEDI9197	TLR7/8	Durvalumab (anti-PD-L1)	Solid tumors	02556463	
	MGN1703	TLR9	Ipilimumab (anti-CTLA-4)	Solid tumors	02668770	
	GLA-SE	TLR4	Radiotherapy	Sarcoma	02180698	
	SD-101		TLR9	Ibrutinib, radiotherapy	Follicular lymphoma	02927964
				Pembrolizumab (anti-PD-1)	Melanoma	02521870
	GSK1795091	TLR4	-	Solid tumors	02798978	
	Poly ICLC	TLR3	Durvalumab (anti-PD-L1) + tremelimumab (anti-CTLA-4)	Solid tumors	02643303	
	G100	TLR4	Pembrolizumab (anti-PD-1)	Solid tumors	02501473	
	MIW815 (ADU-S100)	STING	PDR001 (anti-PD-L1)	Solid tumors	03172936	
	BO-112	MDA5	Anti-PD-1	Solid tumors	02828098	
Epigenetic	Azacitidine + entinostat	DNMT HDAC	Nivolumab (anti-PD-1)	NSCLC	01928576	
	Azacitidine	DNMT	Pembrolizumab (anti-PD-1)	Melanoma	02816021	
	Decitabine	DNMT	MBG453 (TRIM3) PDR001 (anti-PD-L1)	MDS, Leukemia	03066648	
	Decitabine	DNMT	Nivolumab (anti-PD-1)	MDS	03259516	
	Decitabine	DNMT	Ipilimumab (anti-CTLA-4)	MDS, Leukemia	02890329	
	Guadecitabine	DNMT	Durvalumab (anti-PD-L1) + tremelimumab (anti-CTLA-4)	SCLC	03085849	

CTLA-4, cytotoxic T lymphocyte-associated molecule-4; DNMT, DNA methyltransferase; HDAC, histone deacetylase; MDA, melanoma differentiation-associated protein; MDS, myelodysplastic syndromes; NCT, National Clinical Trials; NSCLC, non-small cell lung cancer; PD-1, programmed-cell death 1; PRR, pathogen recognition receptor; SCLC, small cell lung cancer; STING, stimulator of interferon genes complex; TLR, Toll-like receptor.

are expressed in many tissues and play a prominent role in myeloid cells, fibroblasts, hepatocytes and central nervous system cells. While highly expressed in plasmacytoid DCs, RLRs are not essential for IFN type I production by these cells.⁴¹

The synthetic dsRNA BO-112 that activates melanoma differentiation-associated protein 5 (MDA5) demonstrated tumor-specific immune responses with a good toxicity profile in a first in-human trial.⁴² A phase I clinical trial testing the combination of BO-112 with anti-PD-1 antibodies is ongoing in several centers of Spain (Table 1). Other interesting antitumoral compounds in development are the synthetic analogs of dsRNA

polyinosinic:polycytidylic acid (poly I:C) that activate TLR3^{37,43–45} and RIG1/MDA5⁴⁶ and 5' triphosphate small interfering RNA (ppp-siRNA) that activates RIG1 and silences specific oncogenes like BCL-2 *via* RNA-interference. Both types of compounds have demonstrated antitumor activities *in vivo*.⁴⁷

The detection of single stranded RNA (ssRNA) is due to TLR7 and TLR8 which activate NF- κ B, IRF-3 and IRF-7, and lead to the expression of type I IFN, TNF- α , interleukin (IL)-1 and IL-12⁴⁸ (Figure 2). TLR7 is mainly expressed in plasmacytoid DCs and B-cells, while TLR8 is expressed in myeloid DCs and Tregs. Imidazoquinolinamin derivatives, such as imiquimod, approved for the

treatment of basal cell carcinoma, show antitumor effects through TLR7 and TLR8 activation.⁴⁹ Intratumoral resiquimod (R848), a ligand of TLR7 and TLR8, in combination with anti-OX40 in animal models induces a systemic antitumor effect.⁵⁰ MEDI9197 (formerly 3M-052), is a novel TLR7/8 dual agonist formulated for intratumoral injection that has been studied in patients with cutaneous tumors, demonstrating safety and immunogenicity.⁵¹

TLR9 recognizes unmethylated CpG motifs from bacteria and viruses⁵² and RNA:DNA hybrids.⁵³ Its triggering activates plasmacytoid DCs and B-cells.⁵⁴ Several synthetic CpG oligonucleotides have been developed as TLR9 agonists mimicking natural CpG motifs (Table 1). The combination of intratumoral CpG SD-101 with anti-OX40, which triggers a T-cell immune response, demonstrated complete responses and long-term survival in animal models.⁵⁰ Similarly, data from a phase I clinical trial in advanced melanoma showed objective responses in four of the five patients treated with the combination of intratumoral CpG SD-101 and the anti-PD-1 antibody pembrolizumab.⁵⁵

The CDSs cGAS and IFI16 operate *via* stimulator of interferon genes complex (STING) to detect free DNA in the cytosol. When cytosolic DNA is detected by cGAS it catalyzes the STING ligand cGAMP.⁵⁶ STING is then activated, translocates to perinuclear endosomes and recruits TBK1 and IRF3. These molecules are phosphorylated and translocate to the nucleus with subsequent transcription of type I IFNs⁵⁶ (Figure 2). STING is essential for the production of type I IFN signaling in DCs which enables them to present tumor antigens and prime CD8+ T lymphocytes leading to T-cell infiltration into the tumor.^{18,57} STING is located at the cytosolic site of the endoplasmic reticulum membrane⁵⁶ and is activated by DNA from damaged tumor cells that reach the cytosol of DCs.⁵⁸ Some intratumoral-injected compounds can activate STING in mice, like 5,6-dimethylxanthone-4-acetic acid (DMXAA).⁵⁹ In humans, the STING agonist ML RR-S2 CDA (MIW815, ADU-S100) is in clinical development⁶⁰ (Table 1).

Interferons

IFNs were discovered in the late 1950s during replication studies of influenza virus.⁶¹ They are cytokines that play a critical role in innate immune responses against viral infections and

participate in the activation of the adaptive immune response.²⁶ Several types of IFNs have been described including type I IFNs (various IFN- α , IFN- β , and others), the type II IFN γ , and the more recently classified type III IFNs (IFN- λ). Type I IFNs are produced by most cell types: fibroblasts, DCs and hepatocytes (through RIG-1), plasmacytoid DCs (through TLR9 and TLR7/8) and macrophages and hepatocytes (through TLR3 and TLR4). Type II IFNs are mainly secreted by T-helper-1 (Th-1) lymphocytes, CD8+ lymphocytes, NK and NK T-cells.⁶²

All the different types of IFNs signal through IFN receptors that are differently distributed among cell types. IFN- β binds to the IFN- α/β receptor (IFNAR) to further activate production of more type I IFN⁶³ (Figure 3). IFNAR activates janus kinase (JAK) family members JAK1 and Tyk-2, and subsequently signal transducer and activator of transcription 1 (STAT1) and 2 (STAT2). STAT1/2 bind to IRF9 (p48) and form the IFN-stimulated gene factor 3 (ISGF3; Figure 3). ISGF3 initiates the transcription of several interferon-stimulated genes (ISGs) by binding to the promoter region of IFN-stimulated response elements (ISRE; Figure 3). ISGs include PKR, IRF-1 and IRF7. When activated by TBK-1/IKKe, ISGs regulate IFN α gene transcription.⁶⁴ ISGs activate antimicrobial programs that both degrade viral proteins and inhibit cancer cell proliferation⁶⁵ (Figure 3). Type I IFNs also stimulate the adaptive immune response. Specifically, they promote major histocompatibility complex (MHC) class I and II expression on antigen-presenting cells like DCs that is required for efficient T-cell stimulation. Mature DCs are then able to initiate the adaptive immune response by activating antigen-specific naïve T-cells to proliferate and produce type II IFN.^{66,67} The type II IFN receptor is called IFNGR. It activates genes containing a gamma-activated sequence (GAS) through JAK1/2 and STAT1 signaling (Figure 3). Type II IFNs stimulate the adaptive immune response and activate macrophages and NK cells.

The expression levels of ISGs are predictive of the response to immune checkpoint inhibitors in melanoma patients treated with anti-CTLA-4 antibodies.⁶⁸ Likewise, tumor samples from melanoma, ovarian, lung, breast or colorectal cancer can be classified according to high or low expression levels of IFN-stimulated viral defense genes for example IRF7, RIG1, STAT1, IFNB1,




Figure 3. Interferon pathway. Type I IFN binds to the IFNAR. IFNAR activates the JAK family members JAK1 and Tyk-2, with subsequent phosphorylation of signal transducer and activator of transcription 1 (STAT1) and 2 (STAT2) proteins. These proteins form the complex called ISGF3 when they bind to IRF9 (p48). ISGF3 initiates transcription of several ISGs by binding to ISREs in their promoter regions. The receptor of type II IFN is called IFNGR and also initiates induction of JAK1 and JAK2 recruitment with STAT1 homodimers that activate genes containing a GAS. GAS, gamma-activated sequence; IFN, interferon; IFNAR, IFN- α/β receptor; ISG, interferon-stimulated gene; ISGF3, IFN-stimulated gene factor 3; ISRE, IFN-stimulated response element; JAK, Janus kinase.

IFI6 induced by the hypomethylating agent (HMA) 5-aza-cytidine (5-aza).⁶⁹ Our own work shows that lung cancer and melanoma patients with high tumoral expression of IFN- γ have a better outcome with immunotherapy compared with patients with low IFN- γ expression.⁷⁰

The role of DNA methylation in innate immune responses

Epigenetic modulator drugs can restore immunogenicity and immune recognition of tumors. Therefore, there is an increasing interest in combined epigenetic therapy and immunotherapy.⁷¹

DNA methyltransferases (DNMTs) are important players in epigenetic modulation of the innate immune response. In non-small cell lung cancer with mesenchymal phenotype, STAT3 activates DNMT1, which methylates the promoter regions of RIG1 as well as IRF1, immunoproteasomes (PSMB8, PSMB9) and HLA molecules leading

to the reduction of their expression.⁷² STAT3 also inhibits STAT1 expression, a key regulator of the antigen-presentation machinery in epithelial cells.^{70,72}

HMA such as the nucleoside analogs of cytidine, 5-aza and 5-aza-2-deoxycytidine (5-aza-2dc or decitabine) inhibit DNMTs and are currently approved for the treatment of hematologic malignancies.^{73,74} HMA activate the innate immune response through PRRs, but they also have several other activities that make them a good partner to combine with immune checkpoint inhibitors. For instance, HMA reactivate silenced tumor suppressor genes that encode proteins which limit the proliferative and survival capacity of a cell.^{75,76} HMA induce T-cell responses by stimulating HLA I expression. For example, in brain tumors, decitabine promotes the surface presentation of tumor-associated peptides in the context of HLA I and is thus a good candidate to be combined with other

immunotherapeutic regimen.⁷⁷ Furthermore, treatment of tumor cell lines with the DNA methyltransferase inhibitor 5-azacytidine (AZA) enrich tumors in genes involved in immunomodulatory pathways, defining an ‘AZA Immune gene set (AIMs)’ that can classify primary tumors into ‘high’ and ‘low’ AIM gene expression subsets (see Table 1, in Li and colleagues⁷⁸). HMAs induce the expression of chemokines that ultimately re-educate tumor cells to become more immunogenic.^{71,79}

Last but not least, HMAs have antitumor activities by causing the re-expression of endogenous retroviruses (ERVs) in preclinical cancer models. Human ERVs (HERVs) are retroviral elements that have been fixed within the human genome through evolution. About 8% of the human genome is composed of these HERVs, most of which are non-functional due to the accumulation of mutations and epigenetic control. Chiappinelli and colleagues showed in preclinical models that HMAs through ERV expression induce viral mimicry and IFN signaling to increase tumor immunogenicity and recognition.¹³ Other investigators have reported similar findings in colon cancer, demonstrating an IFN response through activation of MDA5 by dsRNA.⁸⁰ This effect, through induction of HERVs, was synergistic with the effect of anti-CTLA-4 or anti-PD-1/PD-L1 antibodies.^{79,13,81} One way to activate TLRs is through induction of dsRNA from HERVs using low doses of HMAs. HERV-derived viral transcripts then increase within the cytosol leading to RIG1, MDA5 and TLR3 activation and subsequently type I IFN production.⁸² The HMA-induced upregulation of HERVs is synergistic with anti-PD-1/PD-L1 antibodies⁸⁰ and anti-CTLA-4 antibodies.^{82,83}

In a subset of non-small cell lung cancer patients, 20% of durable responses were observed when the anti-PD-1 antibody was given after tumor progression under low dose of HMAs, indicating that HMAs may prime tumors for a subsequent response to immunotherapy.^{79,84} Based on the above and other evidences, there is currently a long list of clinical trials in several types of tumors combining immune checkpoint inhibitors with epigenetic drugs (Table 1 and Table 1 in Dunn and colleagues⁷¹).

Interestingly, apart from HMAs, cyclin-dependent kinase 4/6 (CDK4/6) inhibitors also suppress

DNMT1 and induce viral mimicry. The combination of immune checkpoint inhibitors with CDK4/6 inhibitors was found to be synergistic *in vitro* and *in vivo*^{85,86} and clinical trials are now ongoing with this combination [ClinicalTrials.gov identifiers: NCT02791334, NCT02079636, and NCT02779751].

Other factors in the innate immune response and antitumor therapy

NK cells play an important role against tumor cells. Cells infected by viruses, as well as cancer cells, may downregulate surface MHC class I molecules in order to avoid recognition by CD8⁺ T-cells.⁸⁷ Paradoxically, this phenomenon makes them susceptible to NK cells, that are activated when their membrane receptors, killer-cell immunoglobulin-like receptors (KIRs), are not bound to MHC class I molecules. Some cancer cells bypass NK control because they downregulate specific MHC class I molecules, that are needed to present peptides to CD8⁺ T-cells, while they still express MHC class I peptides that serve as KIR ligands.⁸⁸

Viral restriction factors comprise a group of several proteins expressed by cells to suppress viral replication. These restriction factors constitute an early defense against viral infections, and are partly induced by IFNs.⁸⁹ The best characterized restriction factors are: apolipoprotein B messenger RNA editing enzyme catalytic polypeptide-like3 (APOBEC3) proteins, tripartite-motif-containing 5a (TRIM5a), SAM domain and HD domain-containing protein 1 (SAMHD1), Schlafen 11 (SLFN11) and Tetherin.⁸⁹ Their role in the immune response to cancer cells has not been studied, however it is known that some of them are involved in the process of somatic mutagenesis during tumor development^{90,91} APOBEC3 alterations (mutations or overexpression) in cancer cells have been linked to a specific hypermutation status named ‘kataegis’ that correlates with responsiveness to immunotherapy.⁹² Finally, a strong correlation has been reported among APOBEC3 proteins, IFN γ and PD-L1 expression in cancer cells.⁹²

There are several compounds in development with the aim to activate an innate immune response and direct CD8⁺ T-cells into tumors (Table 1). Such strategies include the combination of anti-PD-1 antibodies with oncolytic virotherapy which has shown impressive results in

early clinical trials.⁹³ Melanoma patients treated with the combination of the anti-PD-1 antibody pembrolizumab plus talimogene laherparepvec (T-VEC) had an objective response rate of 62% with 32% of complete responses.¹⁰ Although the exact mechanism by which oncolytic viruses increase the activity of immune checkpoint blockade is unknown, biopsies taken from patients before treatment and after 6 weeks, demonstrated an increase in CD8+ T-cell infiltration as well as increased expression of IFN- γ suggesting a role of innate immune response activation by T-VEC.¹⁰ Similarly, a hybrid of an oncolytic, nonpathogenic poliovirus (PV) and a human rhinovirus (PVSRIPO) demonstrated an increased release of DAMPs through tumor lysis activating a type I IFN response in DCs in different cancer models.⁹

Conclusions and future perspectives

Innate immune responses in the form of IFN production and their subsequent effects provide the appropriate microenvironment for the efficient stimulation of adaptive responses that are key in fighting microbial infections. As tumors are derived from tissue cells that carry mainly self-antigens, they are expected to be purely immunogenic, and may lack innate immune activation, at least initially. Subsequently, the growing tumor can exploit all of the immune suppressive mechanisms that are well described in chronic infections.

Recent clinical trials and preclinical cancer models now impressively demonstrate that stimuli of innate immunity in combination with other immunotherapeutic regimens can significantly augment tumor-specific responses that translate into increased response and cure rates (Figure 1). Thus, suppressive mechanisms and low immunogenicity may be overcome at least when sufficient numbers of tumor neoantigens are present in the tissue.

While presently the PRRs agonists and HMAs stand out, other drugs and drug combinations are under study in numerous trials (Table 1). Furthermore, there are now ongoing trials that include cancer patients with persistent virus infections that have previously been excluded. For example, we are currently exploring several above-mentioned factors and their relationship with innate immune responses in a phase II Spanish Lung Cancer Group clinical trial. The

anti-PD-L1 inhibitor durvalumab is given to HIV-1-infected patients with solid tumors [ClinicalTrials.gov identifier: NCT03094286]. The trial enables us to investigate the effect of immune checkpoint inhibition in reversing cancer pathways and to characterize HIV-specific T-cell functions during persistent HIV infection. The study results of this trial and of others are eagerly awaited.

It seems that after many years of stagnation, we finally face most fruitful and exciting times in tumor immunology.

Funding

AM is supported by a grant from the Spanish Ministry of Economy, Industry and Competitiveness and FEDER grant no. SAF2016-75505-R (AEI/MINEICO/FEDER, UE), and the 'María de Maeztu' Programme for Units of Excellence in R&D (MDM-2014-0370).

Work in Dr Rosell's laboratory is partially supported by a grant from La Caixa Foundation, and by a European Grant (ELBA no 765492).

Conflict of interest statement

The authors declare that there is no conflict of interest.

References

1. Ribas A and Wolchok JD. Cancer immunotherapy using checkpoint blockade. *Science* 2018; 359: 1350–1355.
2. Robert C, Schachter J, Long GV, *et al.* Pembrolizumab versus ipilimumab in advanced melanoma. *N Engl J Med* 2015; 372: 2521–2532.
3. Reck M, Rodriguez-Abreu D, Robinson AG, *et al.* Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N Engl J Med* 2016; 375: 1823–1833.
4. Motzer RJ, Escudier B, McDermott DF, *et al.* Nivolumab versus everolimus in advanced renal-cell carcinoma. *N Engl J Med* 2015; 373: 1803–1813.
5. Tumei PC, Harview CL, Yearley JH, *et al.* PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature* 2014; 515: 568–571.
6. Murphy K and Weaver C. *Janevay's immunobiology*. 9th ed. New York, NY: Garland Science, 2016.

7. Colunga A, Pulliam T and Nghiem P. Merkel cell carcinoma in the age of immunotherapy: facts and hopes. *Clin Cancer Res* 2018; 24: 2035–2043.
8. Seiwert TY, Burtneß B, Mehra R, *et al.* Safety and clinical activity of pembrolizumab for treatment of recurrent or metastatic squamous cell carcinoma of the head and neck (KEYNOTE-012): an open-label, multicentre, phase 1b trial. *Lancet Oncol* 2016; 17: 956–965.
9. Brown MC, Holl EK, Boczkowski D, *et al.* Cancer immunotherapy with recombinant poliovirus induces IFN-dominant activation of dendritic cells and tumor antigen-specific CTLs. *Sci Transl Med* 2017; 9: eaan4220.
10. Ribas A, Dummer R, Puzanov I, *et al.* Oncolytic virotherapy promotes intratumoral T cell infiltration and improves anti-PD-1 immunotherapy. *Cell* 2017; 170: 1109–1119. e10.
11. Chiappinelli KB, Strissel PL, Desrichard A, *et al.* Inhibiting DNA methylation causes an interferon response in cancer via dsRNA including endogenous retroviruses. *Cell* 2017; 169: 361.
12. Gonzalez-Cao M, Iduma P, Karachaliou N, *et al.* Human endogenous retroviruses and cancer. *Cancer Biol Med* 2016; 13: 483–488.
13. Chiappinelli KB, Strissel PL, Desrichard A, *et al.* Inhibiting DNA methylation causes an interferon response in cancer via dsRNA including endogenous retroviruses. *Cell* 2015; 162: 974–986.
14. Balermipas P, Martin D, Wieland U, *et al.* Human papilloma virus load and PD-1/PD-L1, CD8(+) and FOXP3 in anal cancer patients treated with chemoradiotherapy: rationale for immunotherapy. *Oncotarget* 2017; 6: e1288331.
15. Solomon B, Young RJ, Bressel M, *et al.* Prognostic significance of PD-L1+ and CD8+ immune cells in HPV+ oropharyngeal squamous cell carcinoma. *Cancer Immunol Res* 2018; 6: 295–304.
16. Oguejiofor K, Galletta-Williams H, Dovedi SJ, *et al.* Distinct patterns of infiltrating CD8+ T cells in HPV+ and CD68 macrophages in HPV- oropharyngeal squamous cell carcinomas are associated with better clinical outcome but PD-L1 expression is not prognostic. *Oncotarget* 2017; 8: 14416–14427.
17. Fuertes MB, Woo SR, Burnett B, *et al.* Type I interferon response and innate immune sensing of cancer. *Trends Immunol* 2013; 34: 67–73.
18. Spranger S, Dai D, Horton B, *et al.* Tumor-residing Batf3 dendritic cells are required for effector T cell trafficking and adoptive T cell therapy. *Cancer Cell* 2017; 31: 711–723.e4.
19. Dear AE. Epigenetic modulators and the new immunotherapies. *N Engl J Med* 2016; 374: 684–686.
20. Fuertes MB, Kacha AK, Kline J, *et al.* Host type I IFN signals are required for antitumor CD8+ T cell responses through CD8{alpha}+ dendritic cells. *J Exp Med* 2011; 208: 2005–2016.
21. Thompson MR, Kaminski JJ, Kurt-Jones EA, *et al.* Pattern recognition receptors and the innate immune response to viral infection. *Viruses* 2011; 3: 920–940.
22. Seth RB, Sun L and Chen ZJ. Antiviral innate immunity pathways. *Cell Res* 2006; 16: 141–147.
23. Barton GM and Medzhitov R. Toll-like receptors and their ligands. *Curr Top Microbiol Immunol* 2002; 270: 81–92.
24. Altfeld M and Gale M, Jr. Innate immunity against HIV-1 infection. *Nat Immunol* 2015; 16: 554–562.
25. Porta C, Hadj-Slimane R, Nejmeddine M, *et al.* Interferons alpha and gamma induce p53-dependent and p53-independent apoptosis, respectively. *Oncogene* 2005; 24: 605–615.
26. Janeway CA, Jr. and Medzhitov R. Innate immune recognition. *Annu Rev Immunol* 2002; 20: 197–216.
27. Blasius AL and Beutler B. Intracellular toll-like receptors. *Immunity* 2010; 32: 305–315.
28. Xiao T. Innate immune recognition of nucleic acids. *Immunol Res* 2009; 43: 98–108.
29. Roers A, Hiller B and Hornung V. Recognition of endogenous nucleic acids by the innate immune system. *Immunity* 2016; 44: 739–754.
30. Poltorak A, He X, Smirnova I, *et al.* Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science* 1998; 282: 2085–2088.
31. Kang JY, Nan X, Jin MS, *et al.* Recognition of lipopeptide patterns by Toll-like receptor 2-Toll-like receptor 6 heterodimer. *Immunity* 2009; 31: 873–884.
32. Ran S. The role of TLR4 in chemotherapy-driven metastasis. *Cancer Res* 2015; 75: 2405–2410.
33. Huang L, Xu H and Peng G. TLR-mediated metabolic reprogramming in the tumor microenvironment: potential novel strategies for cancer immunotherapy. *Cell Mol Immunol* 2018; 15: 428–437.
34. Onier N, Hilpert S, Arnould L, *et al.* Cure of colon cancer metastasis in rats with the new lipid A OM 174. Apoptosis of tumor cells and immunization of rats. *Clin Exp Metastasis* 1999; 17: 299–306.

35. Okamoto M, Oshikawa T, Tano T, *et al.* Mechanism of anticancer host response induced by OK-432, a streptococcal preparation, mediated by phagocytosis and Toll-like receptor 4 signaling. *J Immunother* 2006; 29: 78–86.
36. Akeda T, Yamanaka K, Kitagawa H, *et al.* Intratumoral injection of OK-432 suppresses metastatic squamous cell carcinoma lesion inducing interferon-gamma and tumour necrosis factor-alpha. *Clin Exp Dermatol* 2012; 37: 193–194.
37. Awasthi S. Toll-like receptor-4 modulation for cancer immunotherapy. *Front Immunol* 2014; 5: 328.
38. Galluzzi L, Vacchelli E, Eggermont A, *et al.* Trial watch: experimental Toll-like receptor agonists for cancer therapy. *Oncoimmunology* 2012; 1: 699–716.
39. Jiang Z, Mak TW, Sen G, *et al.* Toll-like receptor 3-mediated activation of NF-kappaB and IRF3 diverges at Toll-IL-1 receptor domain-containing adapter inducing IFN-beta. *Proc Natl Acad Sci USA* 2004; 101: 3533–3538.
40. Matsumiya T and Stafforini DM. Function and regulation of retinoic acid-inducible gene-I. *Crit Rev Immunol* 2010; 30: 489–513.
41. Loo YM and Gale M, Jr. Immune signaling by RIG-I-like receptors. *Immunity* 2011; 34: 680–692.
42. Marquez Rodas I, Rodriguez-Ruiz ME, Lopez-Tarruella S, *et al.* First-in-human clinical trial with intratumoral BO-112 in solid malignancies: a novel immunotherapy based in double-stranded RNA (dsRNA). *J Clin Oncol* 2017; 35(15 Suppl.): 3082.
43. Amos SM, Pegram HJ, Westwood JA, *et al.* Adoptive immunotherapy combined with intratumoral TLR agonist delivery eradicates established melanoma in mice. *Cancer Immunol Immunother* 2011; 60: 671–683.
44. Chicoine MR, Won EK and Zahner MC. Intratumoral injection of lipopolysaccharide causes regression of subcutaneously implanted mouse glioblastoma multiforme. *Neurosurgery* 2001; 48: 607–614; discussion 14–5.
45. Brody JD, Ai WZ, Czerwinski DK, *et al.* In situ vaccination with a TLR9 agonist induces systemic lymphoma regression: a phase I/II study. *J Clin Oncol* 2010; 28: 4324–4332.
46. Kubler K, the Pesch C, Gehrke N, *et al.* Immunogenic cell death of human ovarian cancer cells induced by cytosolic poly(I:C) leads to myeloid cell maturation and activates NK cells. *Eur J Immunol* 2011; 41: 3028–3039.
47. Poeck H, Besch R, Maihoefer C, *et al.* 5'-Triphosphate-siRNA: turning gene silencing and Rig-I activation against melanoma. *Nat Med* 2008; 14: 1256–1263.
48. Diebold SS. Recognition of viral single-stranded RNA by Toll-like receptors. *Adv Drug Deliv Rev* 2008; 60: 813–823.
49. Li ZJ, Sohn KC, Choi DK, *et al.* Roles of TLR7 in activation of NF-kappaB signaling of keratinocytes by imiquimod. *PLoS One* 2013; 8: e77159.
50. Sagiv-Barfi I, Czerwinski DK, Levy S, *et al.* Eradication of spontaneous malignancy by local immunotherapy. *Sci Transl Med* 2018; 10: eaan4488.
51. Gupta S, Grilley-Olson J, Hong D, *et al.* Safety and pharmacodynamic activity of MEDI9197, a TLR 7/8 agonist, administered intratumorally in subjects with solid tumors. *Cancer Res* 2017; 77: abstract CT091.
52. Ashkar AA and Rosenthal KL. Toll-like receptor 9, CpG DNA and innate immunity. *Curr Mol Med* 2002; 2: 545–556.
53. Rigby RE, Webb LM, Mackenzie KJ, *et al.* RNA: DNA hybrids are a novel molecular pattern sensed by TLR9. *Embo J* 2014; 33: 542–558.
54. Bafica A, Scanga CA, Feng CG, *et al.* TLR9 regulates Th1 responses and cooperates with TLR2 in mediating optimal resistance to Mycobacterium tuberculosis. *J Exp Med* 2005; 202: 1715–1724.
55. Ribas A, Gonzalez R and Kummar S. Phase 1b/2, Open-label, multicenter, dose escalation and expansion trial of intratumoral SD-101 in combination with pembrolizumab in patients with metastatic melanoma. *SMR Meeting* 2016.
56. Woo SR, Fuertes MB, Corrales L, *et al.* STING-dependent cytosolic DNA sensing mediates innate immune recognition of immunogenic tumors. *Immunity* 2014; 41: 830–842.
57. Spaulding E, Fooksman D, Moore JM, *et al.* STING-licensed macrophages prime type I IFN production by plasmacytoid dendritic cells in the bone marrow during severe plasmodium yoelii malaria. *PLoS Pathogens* 2016; 12: e1005975.
58. Corrales L and Gajewski TF. Molecular pathways: targeting the stimulator of interferon genes (STING) in the immunotherapy of cancer. *Clin Cancer Res* 2015; 21: 4774–4779.

59. Corrales L, Glickman LH, McWhirter SM, *et al.* Direct activation of STING in the tumor microenvironment leads to potent and systemic tumor regression and immunity. *Cell Rep* 2015; 11: 1018–1030.
60. Corrales L and Gajewski TF. Endogenous and pharmacologic targeting of the STING pathway in cancer immunotherapy. *Cytokine* 2016; 77: 245–247.
61. Isaacs A and Lindenmann J. Virus interference. I. The interferon. *Proc R Soc Lond B Biol Sci* 1957; 147: 258–267.
62. Kambayashi T, Assarsson E, Lukacher AE, *et al.* Memory CD8⁺ T cells provide an early source of IFN-gamma. *J Immunol* 2003; 170: 2399–2408.
63. Honda K, Yanai H, Takaoka A, *et al.* Regulation of the type I IFN induction: a current view. *Int Immunol* 2005; 17: 1367–1378.
64. Honda K, Yanai H, Negishi H, *et al.* IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* 2005; 434: 772–777.
65. Zitvogel L, Galluzzi L, Kepp O, *et al.* Type I interferons in anticancer immunity. *Nat Rev Immunol* 2015; 15: 405–414.
66. Biron CA, Nguyen KB, Pien GC, *et al.* Natural killer cells in antiviral defense: function and regulation by innate cytokines. *Annu Rev Immunol* 1999; 17: 189–220.
67. Kadowaki N, Antonenko S, Lau JY, *et al.* Natural interferon alpha/beta-producing cells link innate and adaptive immunity. *J Exp Med* 2000; 192: 219–226.
68. Snyder A, Makarov V, Merghoub T, *et al.* Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N Engl J Med* 2014; 371: 2189–2199.
69. Li H, Chiappinelli KB, Guzzetta AA, *et al.* Immune regulation by low doses of the DNA methyltransferase inhibitor 5-azacytidine in common human epithelial cancers. *Oncotarget* 2014; 5: 587–598.
70. Karachaliou N, Gonzalez-Cao M, Crespo G, *et al.* Interferon gamma, an important marker of response to immune checkpoint blockade in non-small cell lung cancer and melanoma patients. *Ther Adv Med Oncol* 2018; 10: 1758834017749748.
71. Dunn J and Rao S. Epigenetics and immunotherapy: the current state of play. *Mol Immunol* 2017; 87: 227–239.
72. Tripathi SC, Peters HL, Taguchi A, *et al.* Immunoproteasome deficiency is a feature of non-small cell lung cancer with a mesenchymal phenotype and is associated with a poor outcome. *Proc Natl Acad Sci USA* 2016; 113: E1555–E1564.
73. Ley TJ, Ding L, Walter MJ, *et al.* DNMT3A mutations in acute myeloid leukemia. *N Engl J Med* 2010; 363: 2424–2433.
74. Popovic R, Shah MY and Licht JD. Epigenetic therapy of hematological malignancies: where are we now? *Ther Adv Hematol* 2013; 4: 81–91.
75. Kazanets A, Shorstova T, Hilmi K, *et al.* Epigenetic silencing of tumor suppressor genes: paradigms, puzzles, and potential. *Biochim Biophys Acta* 2016; 1865: 275–288.
76. Paz MF, Wei S, Cigudosa JC, *et al.* Genetic unmasking of epigenetically silenced tumor suppressor genes in colon cancer cells deficient in DNA methyltransferases. *Hum Mol Genet* 2003; 12: 2209–2219.
77. Shraibman B, Kadosh DM, Barnea E, *et al.* Human Leukocyte Antigen (HLA) peptides derived from tumor antigens induced by inhibition of DNA methylation for development of drug-facilitated immunotherapy. *Mol Cell Proteomics* 2016; 15: 3058–3070.
78. Dietrich PA, Yang C, Leung HH, *et al.* GPR84 sustains aberrant beta-catenin signaling in leukemic stem cells for maintenance of MLL leukemogenesis. *Blood* 2014; 124: 3284–3294.
79. Wrangle J, Wang W, Koch A, *et al.* Alterations of immune response of non-small cell lung cancer with azacytidine. *Oncotarget* 2013; 4: 2067–2079.
80. Roulois D, Loo Yau H, Singhanian R, *et al.* DNA-demethylating agents target colorectal cancer cells by inducing viral mimicry by endogenous transcripts. *Cell* 2015; 162: 961–973.
81. Liu M, Ohtani H, Zhou W, *et al.* Vitamin C increases viral mimicry induced by 5-aza-2'-deoxycytidine. *Proc Natl Acad Sci USA* 2016; 113: 10238–1044.
82. Nusinzon I and Horvath CM. Positive and negative regulation of the innate antiviral response and beta interferon gene expression by deacetylation. *Mol Cell Biol* 2006; 26: 3106–3113.
83. Wang L, Amoozgar Z, Huang J, *et al.* Decitabine enhances lymphocyte migration and function and synergizes with CTLA-4 blockade in a murine ovarian cancer model. *Cancer Immunol Res* 2015; 3: 1030–1041.
84. Juergens RA, Wrangle J, Vendetti FP, *et al.* Combination epigenetic therapy has efficacy in patients with refractory advanced non-small cell lung cancer. *Cancer Discov* 2011; 1: 598–607.

85. Goel S, DeCristo MJ, Watt AC, *et al.* CDK4/6 inhibition triggers anti-tumour immunity. *Nature* 2017; 548: 471–475.
86. Schaer DA, Beckmann RP, Dempsey JA, *et al.* The CDK4/6 inhibitor abemaciclib induces a T cell inflamed tumor microenvironment and enhances the efficacy of PD-L1 checkpoint blockade. *Cell Rep* 2018; 22: 2978–2994.
87. Jonjic S, Babic M, Polic B, *et al.* Immune evasion of natural killer cells by viruses. *Curr Opin Immunol* 2008; 20: 30–38.
88. Guma M, Angulo A and Lopez-Botet M. NK cell receptors involved in the response to human cytomegalovirus infection. *Curr Top Microbiol Immunol* 2006; 298: 207–223.
89. Simon V, Bloch N and Landau NR. Intrinsic host restrictions to HIV-1 and mechanisms of viral escape. *Nat Immunol* 2015; 16: 546–553.
90. Henderson S and Fenton T. APOBEC3 genes: retroviral restriction factors to cancer drivers. *Trends Mol Med* 2015; 21: 274–284.
91. Caval V, Suspene R, Shapira M, *et al.* A prevalent cancer susceptibility APOBEC3A hybrid allele bearing APOBEC3B 3'UTR enhances chromosomal DNA damage. *Nat Commun* 2014; 5: 5129.
92. Boichard A, Tsigelny IF and Kurzrock R. High expression of PD-1 ligands is associated with kataegis mutational signature and APOBEC3 alterations. *Oncoimmunology* 2017; 6: e1284719.
93. Dummer R, Hoeller C, Gruter IP, *et al.* Combining talimogene laherparepvec with immunotherapies in melanoma and other solid tumors. *Cancer Immunol Immunother* 2017; 66: 683–695.

Visit SAGE journals online
[journals.sagepub.com/
home/tam](http://journals.sagepub.com/home/tam)

 SAGE journals