


Association of angiotensin-converting enzyme insertion/deletion (ACE I/D) and angiotensinogen (AGT M235T) polymorphisms with the risk of obesity in a Tunisian population

Wided Khmlaoui¹, Sounira Mehri¹ , Sonia Hammami^{1,2}, Roberto Elosua³ and Mohamed Hammami¹

Journal of the Renin-Angiotensin-Aldosterone System
April-June 2020: 1–7
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1470320320907820
journals.sagepub.com/home/jra


Abstract

Objective: This study aims to determine whether genetic variants in *ACE I/D* and *AGT M235T* are associated with overweight-obesity and body mass index (BMI) in a Tunisian population.

Methods: We designed an age- and sex-matched case-control study. The height and weight were measured and BMI was calculated. A total of 259 overweight-obese patients and 369 healthy controls were genotyped for the *ACE I/D* and *AGT M235T* genes using polymerase chain reaction and restriction fragment length polymorphism.

Results: *ACE I/D* and *AGT M235T* genes were associated with BMI, waist circumference and overweight-obesity ($p \leq 0.001$). In an additive model, the I and the M alleles in *ACE* and *AGT* variants, respectively, were associated with a lower BMI: -1.45 and -2.29 units, respectively. *ACE I/D* genotypes were associated with dyslipidemia; *AGT M235T* genotypes with dyslipidemia and total cholesterol.

Conclusion: These data suggest that variations in *ACE I/D* and *AGT M235T* affect the risk of overweight-obesity, BMI and dyslipidemia, and could point to a key molecular pathway of metabolic syndrome and its related comorbidities.

Keywords

Overweight, obesity, *ACE*, *AGT*, genotypes, Tunisian population

Date received: 10 September 2019; accepted: 18 January 2020

Introduction

In recent years, obesity has become one of the major health challenges worldwide.¹ Recent data from the World Health Organization showed that obesity almost tripled between 1975 and 2016.² It is linked to multiple medical complications, affecting quality of life.³ Obesity is a multifactorial disease resulting from complex interactions between genetic (interactions of several genes with each other) and environmental factors.⁴

Polymorphisms in several obesity candidate genes have been the subject of intensive research as obesity is highly influenced by genetics, but a limited number of studies have investigated a possible link between obesity and the renin-angiotensin system (RAS).

The *ACE* gene, mapped on chromosome 17q23, is highly polymorphic in the promoter and coding regions, but because

of the strong linkage disequilibrium in this region, the functional variant of this gene has not yet been determined.⁵ The most studied *ACE* gene polymorphism, the 278-bp insertion (allele I) or deletion (allele D) variant in intron 16, is associated with plasma and cellular *ACE* levels.⁶ The polymorphism results in two homozygous genotypes (DD with 190

¹Biochemistry Laboratory, LR12ES05 “Nutrition-Functional Foods and Vascular Health”, Faculty of Medicine, University of Monastir, Tunisia

²Department of Internal Medicine, CHU F. Bourguiba, Tunisia

³Cardiovascular Epidemiology and Genetics, Institut Hospital del Mar d’Investigacions Mèdiques, Spain

Corresponding author:

Dr. Sounira Mehri, Biochemistry Laboratory, LR12ES05 “Nutrition-Functional Foods and Vascular Health”, Faculty of Medicine, University of Monastir, Monastir, Tunisia, Avicenne St., Monastir, 5019, Tunisia. Email: mehrisounira@yahoo.fr



bp and II with 490 bp) and one heterozygous genotype (ID with 490 bp as well as 190 bp). Cellular and plasma ACE levels are higher in homozygous DD than in homozygous II subjects, with intermediate levels among heterozygous ID subjects.⁶ Unfortunately, data concerning the role of the I/D polymorphism in coronary artery disease risk in different populations are conflicting.^{7–10}

The *AGT* gene, localized on chromosome 1q41-qter, encodes AGT. There is only one haplotype block at the *AGT* locus, and all common single nucleotide polymorphisms (SNPs) identified appear to be in complete linkage disequilibrium with the most intensively studied *M235T* polymorphism.¹¹ Although the functional variant has not yet been definitively identified,¹² the *T235* allele has been consistently associated with cardiovascular disease.^{11,13,14}

Recent genome-wide association studies have discovered several loci associated with obesity-related traits.¹⁵ Most of these studies have been performed in European, Asian or Caucasian populations. However, information about the genetic background of obesity in North African populations is scanty.

Our main aim was to determine the effects of the angiotensin-converting enzyme insertion/deletion (*ACE I/D*) and the angiotensinogen (*AGT M235T*) gene polymorphisms on the risk of overweight-obesity and its associations with body mass index (BMI) in a Tunisian population.

Subjects and methods

Design and subjects

An age- and sex-matched case-control study was designed. Cases were individuals with overweight or obesity recruited from the department of endocrinology from Fattouma Bourguiba University Hospital (Monastir, Tunisia). All these patients were prospectively invited to participate in this study. Controls were normal-weight individuals, and were also prospectively selected and invited to participate among those attending a routine checkup as part of annual physical examination. Controls were living in the same geographic area as cases and without any history of hypertension or diabetes. Samples were collected during the period from January 2018 to April 2019.

Patients were individuals who had essential hypertension and were treated with selective antihypertensive medication (diuretics, beta blockers, calcium channel blockers) for about 3 years or more than that in monotherapy or combination therapy.

Anthropometric variables

Weight (kg), height (cm) and waist circumference (WC, cm) were determined according to a standard protocol in the Nutrition Center by trained nurses. BMI was calculated as weight (kg)/(height)(height)m². Participants were categorized

as normal body weight: 18.5–24.9 kg/m², overweight BMI: 25–29.9 kg/m² or obese BMI: >30 kg/m² (National Heart, Lung, and Blood Institute in cooperation with the National Institute of Diabetes and Digestive and Kidney Diseases, 1998).¹⁶

Sample collection and biochemical Markers

After an overnight fasting, from each of the recruited participants, 5 ml peripheral blood sample was withdrawn by venipuncture of the upper limbs and placed into tubes containing EDTA (for DNA extraction and lipid profiling).

Serum was separated for the analysis of total cholesterol (TC), high-density lipoprotein cholesterol (HDLC) and triglyceride (TG) analysis. Fasting blood glucose was measured at the time of blood collection. Biochemical analysis of TC, TG, HDLC was done using Randox kits (spectrophotometer). Low-density lipoprotein cholesterol was calculated using the Friedewald equation.¹⁷

Genotyping

DNA was extracted from blood samples according to the Miller et al. 1988 protocol.¹⁸ Allele-specific polymerase chain reaction (PCR) of *ACE I/D* polymorphism (specific primer sequences, forward oligo: 5'CTGGAG ACCACTCCCATCCTTTCT3' and reverse oligo: 5'GAT GTGGCCATCACATTCGTCAGAT3') was done following a standardized protocol.¹⁹ DD homozygotes were re-amplified using specific primers (forward oligo: 5'TGG GACCACAGCGCCCGCCACTAC3' and reverse oligo: 5'TCGCCAGCCCTCCCATGCCATAA3') in order to avoid misclassification of heterozygotes as homozygotes.²⁰ The *AGT M235T* genotype was determined by PCR amplification (specific primer sequences, forward oligo: 5'CAG GGTGCTGTCCACACTGGACCCC3' and reverse oligo: 5'CCGTTTGTGCAGGGCCTGGCTCTC3') followed by digestion with restriction enzyme *Tth1111* according to the described method.²¹

Ethical considerations

This study was reviewed and approved by our hospital ethical committee. Participants were informed that participation is voluntary, and written consent was obtained from each participant after discussing the objective of the study and before being subjected to the questionnaire. No names were recorded on the questionnaires.

Statistical analysis

Hardy–Weinberg expectation for the genotypic distributions of SNPs was investigated with a Chi Square goodness of fit test with one degree of freedom.

Table 1. Demographic and biochemical characteristics of the participants in the study stratified by the presence of overweight-obesity.

	Case (n=259)	Control (n=302)	p-value
Age (years) *	48.88 (14.69)	48.27 (13.70)	0.612
Sex (M/F)	155/104	161/141	0.125
Smoking, n (%)	174 (67.2)	184 (60.9)	0.135
Hypertension, n (%)	140 (54.1)	0 (0.0)	<0.001
Dyslipidemia, n (%)	150 (57.9)	0 (0.0)	<0.001
Diabetes, n (%)	100 (38.6)	0 (0.0)	<0.001
Glucose, mmol/l *	6.01 (1.31)	4.56 (1.86)	<0.001
Triglycerides, mmol/l *	1.49 (0.83)	1.39 (0.65)	0.113
Total cholesterol, mmol/l *	5.44 (1.69)	4.47 (0.99)	<0.001
HDL-cholesterol, mmol/l *	1.41 (0.52)	1.20 (0.50)	<0.001
LDL-cholesterol, mmol/l *	3.11 (1.39)	2.58 (0.81)	<0.001
BMI, kg/m² *	31.1 (4.37)	21.7 (2.22)	<0.001
Waist circumference, cm *	119.58 (13.06)	98.33 (11.67)	<0.001
Normal weight, n (%)	–	302 (100)	
Overweight, n (%)	102 (39.4)	–	
Obesity, n (%)	157 (60.6)	–	

*Mean (Standard Deviation).

HDL: high density lipoprotein; LDL: low density lipoprotein; BMI: body mass index.

Continuous variables are presented as mean (standard deviation) and categorical as frequency (percentage). We used ANOVA and Chi Square to compare continuous and categorical variables distribution between groups, respectively. Those variables associated with the genetic variants of interest and with BMI or overweight-obesity were considered as potential confounders. We used multivariate logistic regression analysis and multivariate linear regression analysis to evaluate the association between genetic variants and overweight-obesity and BMI, respectively. These multivariate models were adjusted for all the confounder variables. To correct for multiple comparisons we used the Bonferroni correction to establish the statistical significance threshold (p -value=0.05/3 genetic variants=0.017).

A multi-locus genetic risk score (GRS) was computed for each individual as the sum of the number of risk alleles across the variants that were associated with overweight-obesity or BMI.

The statistical analyses were carried out using SPSS 22.0 statistical software package for social sciences (SPSS, Chicago, IL, USA).

Results

The characteristics of the participants in this study are shown in Table 1. Finally, 259 overweight-obese and 302 normal-weight Tunisian subjects were included. The mean age of the participants was 48.88 ± 14.69 years for patients and 48.27 ± 13.70 years for the control group. The characteristics of the two groups were similar in terms of age, sex, smoking, diabetes and TG. Overweight-obese individuals

presented with a higher proportion of hypertension, dyslipidemia, and with higher levels of TC, LDL and HDL cholesterol, BMI and WC.

Genotype distributions of all studied polymorphisms were compatible with Hardy–Weinberg expectation in cases and controls. The distribution of the *AGT* *M235T* and *ACE* *I/D* polymorphisms is shown in Table 2.

The frequencies of the polymorphic variants of *AGT* and *ACE* genes were significantly different between the 259 patients and the 302 controls included in the analyses, respectively (*AGT*, $p < 0.001$ and *ACE*, $p < 0.001$).

Furthermore, the frequencies of *AGT* and *ACE* genes were significantly different between overweight and obese patients, respectively (*AGT* MM/MT/TT, 11.7/35.9/52.4% vs. 18.5/32.5/49%, $p = 0.308$ and *ACE* II/ID/DD, 16.5/36.9/46.6% vs. 20.4/39.5/40.1%, $p = 0.302$).

The characteristics of patients across genotypes are shown in Table 3. *ACE* and *AGT* genotypes were both associated with dyslipidemia, BMI, WC and overweight-obesity. *AGT* *M235T* genotype was associated with TC.

The DD genotype and TT genotype, for *ACE* and *AGT* respectively, were related to higher proportion of hypertension, smoking, diabetes and odds of being overweight-obese. No significant correlation was found between smoking and hypertension in patients ($r = 0.89$; $p = 0.163$).

In Table 4, we present the univariate and multivariate effect size on BMI per each of the I and M alleles carried by an individual across genetic variants and in the multi-locus GRS. The results of the multivariate logistic regression analyses are shown in Table 5. The presence of the I

Table 2. The distribution of the AGT M235T and ACE I/D polymorphisms in cases and controls.

	Cases (n=259)	Controls (n=302)	p-value
ACE I/D polymorphism			
DD, n (%)	111 (42.9)	43 (14.2)	<0.001
ID, n (%)	100 (38.6)	123 (40.7)	
II, n (%)	48 (18.5)	136 (45.0)	
AGT M235T polymorphism			
TT, n (%)	131 (50.6)	56 (18.5)	<0.001
MT, n (%)	88 (34.0)	129 (42.7)	
MM, n (%)	40 (15.4)	117 (38.7)	

Table 3. Demographic and biochemical characteristics of patients across the two genetic variants analyzed and corresponding genotypes.

	ACE genotypes				AGT genotypes			
	DD (n=111)	ID (n=100)	II (n=48)	p	TT (n=131)	MT (n=88)	MM (n=40)	p
Age (years) *	49.95 (14.73)	48.67 (14.52)	46.73 (14.87)	0.439	49.62 (14.63)	46.07 (14.99)	52.37 (13.32)	0.052
Sex (M/F)	62/49	61/39	31/17	0.499	72/59	55/33	27/13	0.251
Smoking, n (%)	74 (66.7)	69 (69.0)	32 (65.3)	0.887	93 (71.0)	55 (62.5)	27 (65.9)	0.412
Hypertension, n (%)	58 (41.4)	52 (37.1)	30 (21.4)	0.516	75 (53.6)	40 (28.6)	25 (17.9)	0.139
Dyslipidemia, n (%)	83 (55.3)	39 (26.0)	28 (18.7)	<0.001	84 (56.0)	41 (27.3)	25 (16.7)	0.033
Diabetes, n (%)	46 (41.4)	39 (39.0)	24 (49.0)	0.506	48 (36.6)	41 (46.6)	20 (48.8)	0.214
Glucose, mmol/l *	6.03 (1.32)	5.87 (1.28)	6.26 (1.3)	0.232	5.88 (1.34)	6.16 (1.30)	6.14 (1.25)	0.259
Triglycerides, mmol/l *	1.40 (0.71)	1.55 (0.96)	1.58 (0.81)	0.317	1.56 (0.95)	1.37 (0.67)	1.55 (0.74)	0.257
Total cholesterol, mmol/l *	5.54 (1.53)	5.20 (1.80)	5.66 (1.72)	0.193	5.69 (1.75)	5.22 (1.54)	5.06 (1.68)	0.039
HDL cholesterol, mmol/l *	1.44 (0.53)	1.41 (0.51)	1.37 (0.55)	0.753	1.40 (0.49)	1.39 (0.57)	1.50 (0.50)	0.514
LDL cholesterol, mmol/l *	3.13 (1.32)	3.00 (1.38)	3.22 (1.54)	0.640	3.28 (1.43)	3.05 (1.50)	2.62 (0.73)	0.027
BMI, kg/m ² *	32.34 (4.93)	30.16 (3.61)	29.93 (3.67)	<0.001	31.86 (4.76)	30.52 (3.67)	29.6 (3.95)	0.005
Waist, cm *	122.5 (13.7)	117.85 (12.36)	115.8 (12.4)	0.004	122.1 (13.7)	117.4 (13.09)	115.5 (9.64)	0.004

*Mean (Standard Deviation).

HDL: high density lipoprotein; LDL: low density lipoprotein; BMI: body mass index.

and M alleles was also associated with lower odds of being overweight-obese.

Discussion

A number of important research papers have extensively searched the link of *ACE I/D* and *AGT M235T* with several diseases, mainly hypertension^{22,10} and diabetes,²³⁻²⁵ often with highly inconsistent findings throughout various ethnic populations worldwide; associations with overweight and obesity, an important risk factor for cardiovascular diseases, have also been reported.^{9,26,27} In this context, *ACE I/D* and *AGT M235T* polymorphisms association studies in the Tunisian population are very limited, with reports linking them to obesity.²⁸

In the current study on this North African population, we report an association between *ACE I/D* and *AGT M235T* genes, and BMI and overweight-obesity.

Several studies have investigated the association between *ACE I/D* and *AGT M235T* and overweight-obesity

in Asian,^{26,29-31} Caucasian^{32,33} and African populations.^{27,28} Our results are in part consistent with these previous findings, and confirm the relevance of these loci in overweight-obesity also in a North African population.

A number of investigations in this sense have generated mixed results. A study performed by Akin et al. evaluated the *ACE* polymorphism's association among obese patients with insulin resistance (IR) and reported that the DD genotype was significantly higher in IR obese individuals than those without IR.²⁹ Mehri et al. reported an association of BMI with *ACE* gene polymorphism.²⁸ These findings were consistent with those found by El-Hazmi and Warsy, who reported an increased DD genotype frequency in overweight and obese Saudi individuals.²⁶ Mao et al. in the same way found a significant association was observed between DD genotype and overweight/obesity risk in overall populations (Africans, Caucasians and Asians, together) and Africans.⁹ Although some studies have reported significant association between *ACE* polymorphism and obesity, others have found no associations. For instance,

Table 4. Univariate and multivariate linear regression analysis between the analyzed genotypes and BMI, and between the summary genetic risk score (GRS) and BMI.

		β	SE	p-value
ACE I/D (I allele)	Univariate	-2.58	0.29	<0.001
	Multivariate	-1.62	0.23	<0.001
AGT M235T (M allele)	Univariate	-2.74	0.29	<0.001
	Multivariate	-1.66	0.23	<0.001
GRS	Univariate	-2.34	0.18	<0.001
	Multivariate	-1.50	0.15	<0.001

Multivariate analysis adjusted for age, sex, diabetes, hypertension, triglycerides, HDL, LDL.

Table 5. Association between the analyzed genotypes and the presence of overweight-obesity, and between the summary genetic risk score (GRS) and the presence of overweight-obesity. Multivariate logistic regression analysis.

			OR	95%CI	p-value
ACE I/D	DD	I			
	ID	I	0.32	[0.20-0.49]	<0.001
	II	I	0.14	[0.08-0.22]	<0.001
AGT M235T	TT	I			
	MT	I	0.29	[0.19-0.44]	<0.001
		MM	0.15	[0.09-0.23]	<0.001
GRS			0.31	[0.23-0.42]	<0.001

Pan et al. found no close relationship between *ACE* and obesity.³⁴ Motawi et al. published similar findings in a study of Egyptian women in which *ACE* polymorphism was not associated with obesity.³⁵ Rizvi et al. showed that *ACE* gene polymorphisms were not associated with BMI.³⁶ Similar, Pacholczyk et al. reported no significant associations between *ACE* gene polymorphisms and extreme obesity.³⁷ Moreover, Lelis et al. found no associations between BMI and WC (overweight/obesity) and *ACE*.²⁷

Data from literature regarding *AGT* polymorphisms association with obesity were scanty. We have investigated *AGT* polymorphism's influence on overweight and obesity. In our study, a significant association between *AGT* genotypes and BMI, WC and the odds of being overweight-obese was observed, confirming previous findings.³¹ Giacchetti et al. reported a correlation between *AGT* expression in adipose tissue of obese patients and BMI in visceral adipose tissue.³⁸ Similarly, Umemura et al. published a close relationship of increased plasma *AGT* levels with BMI and blood pressure in obese individuals.³⁹ However, Lelis et al. were not able to find association between BMI and WC and *AGT*.²⁷ Moreover, Pratl-Larquemin et al. reported no association between the *AGT* variants and fat mass.⁴⁰

The present study has to be interpreted within the context of its limitations. Herein, only the association between

the genetic polymorphisms of *ACE* and *AGT* with overweight-obesity was studied and we did not address the functionality of the variants. There are no measurements of markers of RAS activation available to correlate directly with the genetic polymorphisms investigated here. Among the limitations we have also to mention the limited number of SNPs analyzed in this study. In addition, the sample size of our study population is modest, hampering our statistical power. Therefore, extending the investigation of the studied associations on a larger sample set is warranted to confirm that the present findings would replicate in other groups. However, we did observe and replicate some associations previously reported in other ethnic groups.

Conclusion

In conclusion, this preliminary case-control study indicates that angiotensin-converting enzyme insertion/deletion and angiotensinogen polymorphisms were significantly associated with overweight-obesity, BMI and dyslipidemia in this sample of the Tunisian population.

Further confirmatory studies with a larger sample number would be necessary to support or contradict our results. Moreover, biological investigation including measurement of *ACE* and *AGT* levels and activity could be beneficial to understand the influences of these genes on the development of this cluster of events.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by The Ministry of Higher Education, Scientific Research and Information and Communication Technologies, Tunisia.

ORCID iD

Sounira Mehri  <https://orcid.org/0000-0002-2221-7193>

References

1. Hruby A and Hu FB. The epidemiology of obesity: A big picture. *Pharmacoeconomics* 2015; 33: 673–689.
2. World Health Organization (WHO). *Obesity and overweight*. Geneva: WHO, 2018.
3. Anari R, Amani R, Latifi SM, et al. Association of obesity with hypertension and dyslipidemia in type 2 diabetes mellitus subjects. *Diabetes Metab Syndr* 2017; 11: 37–41.
4. Van der Klaauw AA and Farooqi IS. The hunger genes: Pathways to obesity. *Cell* 2015; 161, 119–132.
5. Soubrier F, Martin S, Alonso A, et al. High-resolution genetic mapping of the *ACE*-linked QTL influencing

- circulating ACE activity. *Eur J Human Genet* 2002; 10: 553–561.
6. Rigat B, Hubert C, Alhenc-Gelas F, et al. An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *J Clin Invest* 1990; 86: 1343–1346.
 7. Mehri S, Baudin B, Mahjoub S, et al. Angiotensin-converting enzyme insertion/deletion gene polymorphism in a Tunisian healthy and acute myocardial infarction population. *Genet Test Mol Biomarkers* 2010; 14: 85–91.
 8. Bhatti GK, Bhatti JS, Vijayvergiya R, et al. Implications of ACE (I/D) gene variants to the genetic susceptibility of coronary artery disease in Asian Indians. *Ind J Clin Biochem* 2017; 32: 163–170.
 9. Mao S and Huang SM. A meta-analysis of the association between angiotensin-converting enzyme insertion/deletion gene polymorphism and the risk of overweight/obesity. *J Renin Angiotensin Aldosterone Syst* 2015; 16: 687–694.
 10. Mengesha HG, Petrucka P, Spence C, et al. 2019. Effects of angiotensin converting enzyme gene polymorphism on hypertension in Africa: A meta-analysis and systematic review. *Plos One* 14.
 11. Inoue I, Nakajima T, Williams CS, et al. A nucleotide substitution in the promoter of human angiotensinogen is associated with essential hypertension and affects basal transcription in vitro. *J Clin Invest* 1997; 99: 1786–1797.
 12. Isordia-Salas I, Santiago-German D, Cerda-Mancillas MC, et al. Gene polymorphisms of angiotensin-converting enzyme and angiotensinogen and risk of idiopathic ischemic stroke. *Gene* 2019; 688: 163–170.
 13. Bonfim-Silva R, Guimaraes LO, Santos JS, et al. Case-control association study of polymorphisms in the angiotensinogen and angiotensin-converting enzyme genes and coronary artery disease and systemic artery hypertension in African-Brazilians and Caucasian-Brazilians. *J Genet* 2016; 95: 63–69.
 14. Zhai C, Cong HL, Zhang H, et al. 2019; M235T polymorphism in the angiotensinogen gene and cardiovascular disease: An updated meta-analysis of 39 case-control comparisons. *Anatol J Cardiol* 2019; 21: 222–232.
 15. Monda KL, Chen GK, Taylor KC, et al. A meta-analysis identifies new loci associated with body mass index in individuals of African ancestry. *Nat Genet* 2013; 45: 690–696.
 16. National Heart, Lung, Blood Institute, National Institute of Diabetes, Digestive, & Kidney Diseases (US). *Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: The evidence report* (No. 98). National Heart, Lung, and Blood Institute, 1998.
 17. Friedewald WT, Levy RI and Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972; 18: 499–502.
 18. Miller S, Dykes D and Polesky H. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988; 16: 1215.
 19. Rigat B, Hubert C, Corvol P, et al. PCR detection of the insertion/deletion polymorphism of the human angiotensin converting enzyme gene (DCP1) (dipeptidyl carboxypeptidase 1). *Nucl Acid Res* 1992; 20: 1433–1433.
 20. Shanmugam V, Sell KW and Saha BK. Mistyping ACE heterozygotes. *PCR Methods Appl* 1993; 3: 120–121.
 21. Russ AP, Maerz W, Ruzicka V, et al. Rapid detection of the hypertension-associated Met235->Thr allele of the human angiotensinogen gene. *Hum Mol Genet* 1993; 2: 609–610.
 22. Sofronova SI, Kirillina MP, Nikolaev VM, et al. Association of ACE gene polymorphism with hypertension and metabolic risk factors among Indigenous People of the Northern Territory of Yakutia. *Int J Biomed* 2019; 9: 102–105.
 23. Mehri S, Koubaa N, Nakbi A, et al. Relationship between genetic polymorphisms of angiotensin-converting enzyme and methylenetetrahydrofolate reductase as risk factors for type 2 diabetes in Tunisian patients. *Clin Biochem* 2010; 43: 259–266.
 24. Wollinger LM, Dal Bosco SM, Rempel C, et al. Role of ACE and AGT gene polymorphisms in genetic susceptibility to diabetes mellitus type 2 in a Brazilian sample. *Genet Mol Res* 2015; 14: 19110–19116.
 25. Qiao Y-C, Wang M, Pan Y-H, et al. The relationship between ACE/AGT gene polymorphisms and the risk of diabetic retinopathy in Chinese patients with type 2 diabetes. *J Renin Angiotensin Aldosterone Syst* 2018; 19: 1470320317752955.
 26. El-Hazmi MAF and Warsy AS. Increased frequency of angiotensin-converting enzyme DD genotype in Saudi overweight and obese patients. *Ann Saudi Med* 2003; 23: 24–27.
 27. Lelis DDF, Pereira AC, Krieger JE, et al. Polymorphisms of the renin-angiotensin system are not associated with overweight and obesity in a general adult population. *Arch Endocrinol Metab* 2019; 63: 402–410.
 28. Mehri S, Mahjoub S, Hammami S, et al. Renin-angiotensin system polymorphisms in relation to hypertension status and obesity in a Tunisian population. *Mol Biol Rep* 2012; 39: 4059–4065.
 29. Akin F, Turgut S, Bastemir M, et al. Angiotensin-converting enzyme gene polymorphism in overweight and obese Turkish patients with insulin resistance. *DNA Cell Biol* 2010; 29: 207–212.
 30. Kim K. Association of angiotensin-converting enzyme insertion/deletion polymorphism with obesity, cardiovascular risk factors and exercise-mediated changes in Korean women. *Eur J Appl Physiol* 2009; 105: 879–887.
 31. Takakura Y, Yoshida T, Yoshioka K, et al. Angiotensinogen gene polymorphism (Met235Thr) influences visceral obesity and insulin resistance in obese Japanese women. *Metab Clin Exp* 2006; 55: 819–824.
 32. Wacker MJ, Godard MP, McCabellm EH, et al. Sex difference in the association of the angiotensin converting enzyme I/D polymorphism and body mass index. *Med Sci Monit* 2008; 14: CR353–CR357.
 33. Fiatal S, Szigethy E, Szeles G, et al. Insertion/deletion polymorphism of angiotensin-1 converting enzyme is associated with metabolic syndrome in Hungarian adults. *J Renin Angiotensin Aldosterone Syst* 2011; 12: 531–538.
 34. Pan YH, Wang M, Huang YM, et al. ACE gene I/D polymorphism and obesity in 1,574 patients with type 2 diabetes mellitus. *Dis Markers* 2016; 2016: 7420540.
 35. Motawi TK, Shaker OG, Shahin NN, et al. Angiotensin-converting enzyme insertion/deletion polymorphism association with obesity and some related disorders in Egyptian

- females: A case-control observational study. *Nutr Metab* 2016; 13: 68.
36. Rizvi S, Raza ST, Siddiqi Z, et al. Association of angiotensin-converting enzyme and glutathione s-transferase gene polymorphisms with body mass index among hypertensive North Indians. *Sultan Qaboos Univ Med J* 2015; 15: e477–e485.
 37. Pacholczyk M, Ferenc T, Kowalski J, et al. Association of angiotensin-converting enzyme and angiotensin ii type i receptor gene polymorphisms with extreme obesity in Polish individuals. *DNA Cell Biol* 2013; 32: 435–442.
 38. Giacchetti G, Faloia E, Sardu C, et al. Gene expression of angiotensinogen in adipose tissue of obese patients. *Int J Obesity* 2000; 24: S142–S143.
 39. Umemura S, Nyui N, Tamura K, et al. Plasma angiotensinogen concentrations in obese patients. *Am J Hypertens* 1997; 10: 629–633.
 40. Prat-Larquemin L, Oppert JM, Clement K, et al. Adipose angiotensinogen secretion, blood pressure, and AGT M235T polymorphism in obese patients. *Obes Res* 2004; 12: 556–561.